

Instruction Manual

SERVA Blue R Powder

Cat. No. 35051

| General | Dye based on Coomassie [®] * Brilliant Blue R-250. The "R" in the name Brilliant Blue R refers to a blue dye with a reddish hue. | |
|--------------------|--|--|
| Application | Staining of proteins in polyacrylamide and agarose gels ¹ | |
| | Detection of proteins in solution and on cellulose acetate plates^{2,3} | |
| | Detection of lipids in TLC ⁴ | |
| | Determination of CMC of detergents ⁵ | |
| Storage | Store powder dry and dark at +15 $^{\circ}$ C - +30 $^{\circ}$ C. If st ored as indicated, shelf life of powder is ca. 3 – 5 years. | |
| Solubility | Maximum solubilities: 70 mg/ml in water, 20 mg/ml in ethyl-ene glycol monomethyl ether (methyl cellosolve), 10 mg/ml in EtOH. | |
| Staining protocol: | | |
| Solutions | Fixing solution | 10 % glacial acetic acid, 40 % EtOH (10 ml glacial acetic acid/40 ml EtOH/50 ml H_2O) |
| | Stock solution I | 0.2 % SERVA Blue R in 90 % EtOH (100 mg in 50 ml 90 % EtOH) Solution is stable at RT for several months. |
| | Stock solution II | 20 % Acetic acid |
| | Destainer I | conforms to fixing solution |
| | Destainer II | 10 % glacial acetic acid, 20 % EtOH |
| Staining | 1. Fix gel for 30 min. in 100 ml fixing solution. | |
| | 2. Prior to use mix 50 ml stock solution I with 50 ml stock solution II, stain gel in solution for 20 min. | |
| Entfärbung | For a first rapid destain, incubate gel for ca. 0.5 - 5 min. (dependent on gel thickness) in destainer I. Destain gel further as long as needed in destainer II (visual inspection). | |
| | 2. Rinse destained gel 2x for 5 min. in H ₂ O dist. | |

¹Wilson, C.M., Anal. Biochem., 96, 263 (1979), ²McNight, G.S., Anal. Biochem., 78, 86 (1977), ³Fazekas, S., et al., Biochim. Biophys. Acta, 71, 377 (1963), ⁴Rosenthal, K.S., et al., Anal. Chem., 55, 1115 (1983).
*Coomassie is a tradename of ICI, UK

Ver. 04/07