

Specifications: Acrylamide-Bis (Powder, Solutions)

Description	A 290 nm	pH	Conductivity (µS)	Content of free acrylic acid	Cat.No.
Acrylamide 4X analytical grade	< 0.6 (5 %)	ca. 6 (5 %)	< 20 (40 %)	< 0.002 %	10674
Acrylamide 2X research grade	< 0.7 (5 %)	ca. 6 (5 %)	< 100 (40 %)	< 0.03 %	10675
Acrylamide 4X molecular biology grade	< 0.6 (5 %)	ca. 6 (5 %)	< 20 (40 %)	< 0.002 %	10678
Acrylamide 4X solution (40 % w/v)	< 0.6 (5 %)	6 – 8 (5 %)	< 100	< 0.03 %	10677
N,N'-Methylene bisacrylamide 2X analytical grade	max. 0.25 (1 %)	5.5 - 7.5 (1 %)			29195
N,N'-Methylene bisacrylamide 2X solution (2 % w/v)	ca. 0.25 (1 %)	6 – 8 (1 %)			29197
Acrylamide/Bis Solution, 19:1 (40 % (w/v), 5 % C)	< 0.7 (5 %)	6 – 8 (5 %)	< 100	< 0.03 %	10679
Acrylamide/Bis Solution, 29:1 (40 % (w/v), 3.3 % C)	< 0.7 (5 %)	6 – 8 (5 %)	< 100	< 0.03 %	10680
Acrylamide/Bis Solution, 37.5:1 (40 % (w/v), 2.6 % C)	< 0.7 (5 %)	6 – 8 (5 %)	< 100	< 0.03 %	10681
Acrylamide/Bis Solution, 29:1 (30 % (w/v), 3.3 % C)	< 0.7 (5 %)	6 – 8 (5 %)	< 100	< 0.03 %	10687
Acrylamide/Bis Solution, 37.5:1 (30 % (w/v), 2.6 % C)	< 0.7 (5 %)	6 – 8 (5 %)	< 100	< 0.03 %	10688